

Quantum Gravity

Enrique Álvarez

*Instituto de Física Teórica UAM/CSIC, C-XVI, and Departamento de Física Teórica,
C-XI,
Universidad Autónoma de Madrid E-28049-Madrid, Spain*¹

Abstract

General lectures on quantum gravity.

1 General Questions on Quantum Gravity

It is not clear at all what is the problem in quantum gravity (cf. [3] or [8] for general reviews, written in the same spirit as the present one). The answers to the following questions are not known, and I believe it can do no harm to think about them before embarking in a more technical discussion.

To begin, it has been proposed that gravity should not be quantized, owing to its special properties as determining the background on which all other fields propagate. There is a whole line of thought on the possibility that gravity is not a fundamental theory, and this is certainly an alternative one has to bear in mind. Indeed, even the *holographic principle* of G. 't Hooft, to be discussed later, can be interpreted in this sense.

¹Lectures given in the 40th Karpacz Winter School.

Granting that, the next question is whether it does make any sense to consider gravitons propagating in some background; that is, whether there is some useful approximation in which there is a particle physics approach to the physics of gravitons as quanta of the gravitational field. A related question is whether semiclassical gravity, i.e., the approximation in which the source of the classical Einstein equations is replaced by the expectation value of the energy momentum tensor of some quantum theory has some physical ([22]) validity in some limit. We shall say more on this problems towards the end.

At any rate, even if it is possible at all, the at first sight easy problem of graviton interactions in an otherwise flat background has withstood analysis of several generations of physicists. The reason is that the coupling constant has mass dimension -1 , so that the structure of the perturbative counterterms involve higher and higher orders in the curvature invariants (powers of the Riemann tensor in all possible independent contractions), schematically,

$$S = \frac{1}{2\kappa_R^2} \int R + \int R^2 + \kappa_R^2 \int R^4 + \dots \quad (1)$$

Nobody knows how to make sense of this approach, except in one case, to be mentioned later on.

It could be possible, *sensu stricto* to stop here. But if we believe that quantum gravity should give answers to such questions as to the fate of the initial cosmological singularity, its is almost unavoidable to speak of the wave function of the universe. This brings its own set of problems, such as to whether it is possible to do quantum mechanics without classical observers or whether the wave function of the Universe has a probabilistic interpretation. Paraphrasing C. Isham [38], one would not known when to qualify a probabilistic prediction on the whole Universe as a successful one.

The aim of the present paper is to discuss in some detail established results on the field. In some strong sense, the review could be finished at once, because there are none. There are, nevertheless, some interesting attempts, which look promising from certain points of view. Perhaps the two approaches that have attracted more attention have been the loop

approach, on the one hand and strings on the other. We shall try to critically assess prospects in both. Interesting related papers are [34][59].

Even if for the time being there is not (by far) consensus on the scientific community of any quantum gravity physical picture, many great physicist have not been able to resist the temptation of working (usually only for a while) on it. This has produced a huge spinoff in quantum field theory; to name only a few, constrained quantization, compensating ghosts, background field expansion and topological theories are concepts or techniques first developed in thinking about these problems, and associated to the names of Dirac, Pauli, Weinberg, Feynman, deWitt, Witten etc. In many cases, more or less surprising relationships have been found with quantum gauge theories. There are probably more in store, if one is to judge from the success of the partial implementation of holographic ideas in Maldacena's conjecture (more on this later).

This should be kept into account when reading the references. We have not attempted to be comprehensive, and we have used only the references familiar to us; but in some of the references, in particular in our own review article of 1989 ([3] there are more entry points into the vast literature. After all, paraphrasing Feynman, we still do not know what could be relevant in a field until the main problems are solved.

2 The issue of background independence

One of the main differences between both attacks to the quantum gravity problem is the issue of background independence, by which it is understood that no particular background should enter into the definition of the theory itself. Any other approach is purportedly at variance with diffeomorphism invariance.

Work in particle physics in the second half of last century led to some understanding of ordinary gauge theories. Can we draw some lessons from there?

Gauge theories can be formulated in the *background field approach*, as introduced by

B. de Witt and others (cf. [20]). In this approach, the quantum field theory depends on a background field, but not on any one in particular, and the theory enjoys background gauge invariance.

Is it enough to have quantum gravity formulated in such a way? ²

It can be argued that the only vacuum expectation value consistent with diffeomorphisms invariance is

$$\langle 0 | g_{\alpha\beta} | 0 \rangle = 0 \tag{2}$$

in which case the answer to the above question ought to be in the negative, because this is a singular background and curvature invariants do not make sense. It all boils down as to whether the ground state of the theory is diffeomorphism invariant or not. There is an example, namely three-dimensional gravity in which invariant quantization can be performed [70]. In this case at least, the ensuing theory is almost topological.

In all attempts of a canonical quantization of the gravitational field, one always ends up with an (constraint) equation corresponding physically to the fact that the total hamiltonian of a parametrization invariant theory should vanish. When expressed in the Schrödinger picture, this equation is often dubbed the *Wheeler-de Witt equation*. This equation is plagued by operator ordering and all other sorts of ambiguities. It is curious to notice that in ordinary quantum field theory there also exists a Schrödinger representation, which came recently to be controlled well enough as to be able to perform lattice computations ([42]).

Gauge theories can be expressed in terms of gauge invariant operators, such as Wilson loops . They obey a complicated set of equations, the loop equations, which close in the large N limit as has been shown by Makeenko and Migdal ([43]). These equations can be properly regularized, e.g. in the lattice. Their explicit solution is one of the outstanding challenges in theoretical physics. Although many conjectures have been advanced in this

²This was, incidentally, the way G. Hooft and M. Veltman did the first complete one-loop calculation ([65]).

direction, no definitive result is available.

3 The canonical approach.

It is widely acknowledged that there is a certain tension between a $(3 + 1)$ decomposition implicit in any canonical approach, privileging a particular notion of time, and the beautiful geometrical structure of general relativity, with its invariance under general coordinate transformations.

Let us now nevertheless explore how far we can go on this road, following the still very much worth reading work of deWitt ([20]).

If we are given a spacelike surface (which will represent physically all spacetime events to which it will be assigned a fixed time), say

$$y^\alpha = f^\alpha(x^i) \tag{3}$$




Figure 1: Spacelike surface of codimension one.

The tangent vectors to the surface are

$$\xi_i \equiv \partial_i f^\alpha \partial_\alpha \tag{4}$$

and the induced metric (that is, the pull-back to the surface of the spacetime metric) is

$$h_{ij} \equiv g_{\alpha\beta} \xi_i^\alpha \xi_j^\beta \tag{5}$$

The unit normal is then defined as;

$$g_{\alpha\beta}n^\alpha\xi_i^\beta = 0$$

$$n^2 \equiv g_{\alpha\beta}n^\alpha n^\beta = 1$$

We are interested now in a set of such surfaces which covers all spacetime; that is, a foliation of (a portion of) the spacetime; namely a one-parameter family of spacelike disjoint hypersurfaces

$$\Sigma_t \equiv \{y^\alpha = f^\alpha(x^i, t)\} \quad (6)$$




Figure 2: ADM lapse and shift variables.

In a classical analysis Arnowitt, Deser and Misner (ADM) ([1]) characterized the embedding via two functions: the *lapse* and the *shift*: we first define the vector

$$N^\alpha \equiv \frac{\partial f^\alpha}{\partial t} \quad (7)$$

in terms of which the lapse, N , is just the projection in the direction of the normal, and the *shift*, N_i the (three) projections tangent to the hypersurface.

$$N^\alpha = N n^\alpha + N^i \xi_i^\alpha \quad (8)$$

All this amounts to a particular splitting of the full spacetime metric:

$$ds^2 = g_{\alpha\beta}dx^\alpha dx^\beta = g_{\alpha\beta}df^\alpha df^\beta = g_{\alpha\beta}(N^\alpha dt + \xi_i^\alpha dx^i)(N^\beta dt + \xi_j^\beta dx^j) = N^2 dt^2 + h_{jk}(N^j dt + dx^j)(N^k dt + dx^k) \quad (9)$$

or, what is the same,

$$g_{\mu\nu} = h^{ij}\xi_{i\mu}\xi_{j\nu} + n_\mu n_\nu \quad (10)$$

All surfaces which are equivalent from the intrinsic point of view, can be however embedded differently; the extrinsic curvature discriminates between them:

$$K_{ij} = -\xi_i^\alpha \nabla_\rho n_\alpha \xi_j^\rho \quad (11)$$

The Gauss- Codazzi equations relate intrinsic curvatures associated with the intrinsic geometry in the hypersurface with spacetime curvatures precisely through the extrinsic curvature:

$$R[h]_{lijk} = R[g]_{\alpha\beta\sigma\rho}\xi_l^\alpha \xi_i^\beta \xi_j^\sigma \xi_k^\rho - K_{ij}K_{lk} + K_{ik}K_{lj} \quad (12)$$

and

$$\nabla[h]_k K_{ij} - \nabla[h]_j K_{ik} = R_{\alpha\beta\sigma\rho} n^\alpha \xi_i^\beta \xi_j^\rho \xi_k^\sigma \quad (13)$$

whereas the curvature scalar is given by

$$R = R^{\alpha\beta}{}_{\alpha\beta} = 2R^{ni}{}_{ni} + R^{ij}{}_{ij} \quad (14)$$

In terms of this splitting, the Einstein-Hilbert action reads:

$$L_{EH} \equiv \sqrt{g}R[g] = N\sqrt{h}(R[h] + K_{ij}K^{ij} - K^2) - \partial_\alpha V^\alpha \quad (15)$$

with

$$V^\alpha = 2\sqrt{g}(n^\beta \nabla_\beta n^\alpha - n^\alpha \nabla_\beta n^\beta) \quad (16)$$

Primary constraints appear when defining canonical momenta:

$$p^\mu \equiv \frac{\partial L}{\partial \dot{N}_\mu} \sim 0 \quad (17)$$

the momenta conjugate to the spatial part of the metric is:

$$\pi^{ij} \equiv \frac{\delta L}{\delta \dot{h}_{ij}} = -h^{1/2}(K^{ij} - Kh^{ij}) \quad (18)$$

The canonical conmutation relations yield:

$$\{\pi^{ij}(\vec{x}), h_{kl}(\vec{x}')\} = -\delta(\vec{x}, \vec{x}') \frac{1}{2}(\delta_k^i \delta_l^j + \delta_k^j \delta_l^i) \quad (19)$$

The total Hamiltonian reads

$$H \equiv \int d^3x (\pi_\mu \dot{N}^\mu + \pi^{ij} \dot{h}_{ij} - L) = \int d^3x (N\mathcal{H} + N^i \mathcal{H}_i) \quad (20)$$

where

$$\mathcal{H}(h, \pi) = h^{-1/2}(\pi_{ij}\pi^{ij} - \frac{1}{2}\pi^2) - h^{1/2}R[h] \quad (21)$$

and

$$\mathcal{H}_i(h, \pi) = -2h_{ik}\partial_j\pi^{kj} - (2\partial_j h_{ki} - \partial_i h_{kj})\pi^{kj} = -2\nabla[h]_j \pi_i{}^j \quad (22)$$

The system of constraints is now consistent (that is, that the classical time evolution of the constraints is still a linear combination of constraints):

$$\dot{p}^\mu = \{p^\mu, H\} = (\mathcal{H}, \mathcal{H}_i) \sim 0 \quad (23)$$

Second class constraints

$$N^\mu = f^\mu \quad (24)$$

can now be imposed. The whole hamiltonian analysis boils down to the two constraint equations

$$\mathcal{H} = 0$$

$$\mathcal{H}_i = 0$$

Much of the preceding analysis is actually quite generic for generally covariant systems. The full set of constraints obeys the Dirac-Schwinger algebra

$$\begin{aligned}
\{\mathcal{H}(\vec{x}), \mathcal{H}(\vec{y})\} &= [\mathcal{H}^i(\vec{x}) + \mathcal{H}^i(\vec{y})] \partial_i \delta(\vec{x}, \vec{y}) \\
\{\mathcal{H}_i(\vec{x}), \mathcal{H}(\vec{y})\} &= \mathcal{H}(\vec{x}) \partial_i \delta(\vec{x}, \vec{y}) \\
\{\mathcal{H}_i(\vec{x}), \mathcal{H}_j(\vec{y})\} &= \mathcal{H}_i(\vec{y}) \partial_j \delta(\vec{x}, \vec{y}) + \mathcal{H}_j(\vec{x}) \partial_i \delta(\vec{x}, \vec{y})
\end{aligned} \tag{25}$$

which is nothing else than the Σ -projected algebra of the $\text{Diff}(\text{M})$ group.

Usually no reduction is made on the dynamical variables of the system, which amounts to keep h_{ij}, π^{ij} as (redundant) quantum variables. It is not clear how singular metrics can be avoided, because it is not easy to impose the condition that the metric is a positive definite operator.

Physical states in the Hilbert space are provisionally defined à la Dirac

$$\begin{aligned}
\hat{\mathcal{H}}|\psi\rangle &= 0 \\
\hat{\mathcal{H}}_i|\psi\rangle &= 0
\end{aligned}$$

It has been realized since long that this whole approach suffers from the *frozen time problem*, i.e., the Hamiltonian reads

$$H \equiv \int d^3x (N\mathcal{H} + N^i\mathcal{H}_i) \tag{26}$$

so that acting on physical states

$$\hat{H}|\psi\rangle = 0 \tag{27}$$

in such a way that Schrödinger's equation

$$i\frac{\partial}{\partial t}|\psi\rangle = \hat{H}|\psi\rangle \tag{28}$$

seemingly forbids any time dependence.

There are many unsolved problems in this approach, which has been kept at a formal level. The first one is an obvious operator ordering ambiguity owing to the nonlinearity. In the same vein, it is not clear whether it is possible to make the constraints hermitian. Besides, it is not clear that one recovers the full Diff invariance from the Dirac- Schwinger algebra. Actually, it is not known whether this is necessary; that is, what is the full symmetry of the quantum theory.

We can proceed further, still formally³, using the Schrödinger representation defined in such a way that

$$(\hat{h}_{ij}\psi)[h] \equiv h_{ij}(x)\psi[h] \quad (29)$$

and

$$(\hat{\pi}^{ij}\psi)[h] \equiv -i\hbar \frac{\delta\psi}{\delta h_{ij}(x)}[h] \quad (30)$$

If we assume that diffeomorphisms act on wave functionals as:

$$\psi[f^*h] = \psi[h] \quad (31)$$

then the whole setup for the quantum dynamics of the gravitational field lies in Wheeler's *superspace* (nothing to do with supersymmetry) which is the set of three-dimensional metrics modulo three-dimensional diffs : $Riem(\Sigma)/Diff(\Sigma)$.

The Hamiltonian constraint then implies the famous *Wheeler-de Witt* equation.

$$-\hbar^2 2\kappa^2 G_{ijkl} \frac{\delta^2\psi}{\delta h_{ik}\delta h_{jl}}[h] - \frac{\hbar}{2\kappa^2} R^{(3)}[h]\psi[h] = 0 \quad (32)$$

where the de Witt tensor is:

$$G_{ijkl} \equiv \frac{1}{\sqrt{h}} \left(h_{ij}h_{kl} - \frac{1}{2}h_{ik}h_{jl} \right) \quad (33)$$

³It is bound to be formal as long as the problem of the infinities is not fully addressed. We know from the analysis of this representation for gauge theories in the lattice that those are the most difficult problems to solve.

Needless to say, this equation, suggestive as it is, is plagued with ambiguities. The manifold of positive definite metrics has been studied by deWitt. He showed that it has signature $(-1, +1^5)$, where the timelike coordinate is given by the breathing mode of the metric:

$$\zeta = \sqrt{\frac{32}{3}} h^{1/4} \quad (34)$$

and in terms of other five coordinates ζ^a orthogonal to the timelike coordinate, the full metric reads

$$ds^2 = -d\zeta^2 + \frac{3}{32} \zeta^2 g_{ab} d\zeta^a d\zeta^b \quad (35)$$

with

$$g_{ab} = \text{tr } h^{-1} \partial_a h h^{-1} \partial_b h \quad (36)$$

The five dimensional submanifold with metric g_{ab} is the coset space

$$SL(3, \mathbb{R})/SO(3) \quad (37)$$

It has been much speculated whether the timelike character of the dilatations lies at the root of the concept of time. The Wheeler-deWitt equation can be written in a form quite similar to the Klein-Gordon equation:

$$\left(-\frac{\partial^2}{\partial \zeta^2} + \frac{32}{3\zeta^2} g^{ab} \partial_a \partial_b + \frac{3}{32} \zeta^2 R^{(3)} \right) \Psi = 0 \quad (38)$$

The analogy goes further in the sense that also here there is a naturally defined scalar product which is not positive definite:

$$(\psi, \chi) \equiv \int_{\Sigma} \psi^* d\Sigma^{ij} G_{ijkl} \frac{\delta \chi}{i \delta h_{kl}} - \chi^* d\Sigma^{ij} G_{ijkl} \frac{\delta \psi}{i \delta h_{kl}} \quad (39)$$

4 Using Ashtekar and related variables

The whole philosophy of this approach is canonical, i.e., an analysis of the evolution of variables defined classically through a foliation of spacetime by a family of spacelike three-surfaces Σ_t . The standard choice in this case as we have just reviewed, is the three-dimensional metric, g_{ij} , and its canonical conjugate, related to the extrinsic curvature.

Here, as in any canonical approach the way one chooses the canonical variables is fundamental.

Ashtekar's clever insight started from the definition of an original set of variables ([10]) stemming from the Einstein-Hilbert lagrangian written in the form ⁴

$$S = \int e^a \wedge e^b \wedge R^{cd} \epsilon_{abcd} \quad (40)$$

where e^a are the one-forms associated to the tetrad,

$$e^a \equiv e^a_\mu dx^\mu. \quad (41)$$

Tetrads are defined up to a local Lorentz transformation

$$(e^a)' \equiv L^a_b(x) e^b \quad (42)$$

The associated $SO(1,3)$ connection one-form ω^a_b is usually called the *spin connection*. Its field strength is the curvature expressed as a two form:

$$R^a_b \equiv d\omega^a_b + \omega^a_c \wedge \omega^c_b. \quad (43)$$

Ashtekar's variables are actually based on the $SU(2)$ self-dual connection

$$A = \omega - i * \omega \quad (44)$$

Its field strength is

$$F \equiv dA + A \wedge A \quad (45)$$

The dynamical variables are then $(A_i, E^i \equiv F^{0i})$. The main virtue of these variables is that constraints are then linearized. One of them is exactly analogous to Gauss'law:

$$D_i E^i = 0. \quad (46)$$

There is another one related to three-dimensional diffeomorphisms invariance,

$$\text{tr } F_{ij} E^i = 0 \quad (47)$$

⁴Boundary terms have to be considered as well. We refer to the references for details.

and, finally, there is the Hamiltonian constraint,

$$tr F_{ij} E^i E^j = 0 \quad (48)$$

On a purely mathematical basis, there is no doubt that Astekhar's variables are of a great ingenuity. As a physical tool to describe the metric of space, they are not real in general. This forces a reality condition to be imposed, which is awkward. For this reason it is usually preferred to use the Barbero-Immirzi ([13][37]) formalism in which the connexion depends on a free parameter, γ ,

$$A_a^i = \omega_a^i + \gamma K_a^i \quad (49)$$

ω being the spin connection and K the extrinsic curvature. When $\gamma = i$ Astekhar's formalism is recovered; for other values of γ the explicit form of the constraints is more complicated. Thiemann ([67]) has proposed a form for the Hamiltonian constraint which seems promising, although it is not clear whether the quantum constraint algebra is isomorphic to the classical algebra (cf.[54]). A comprehensive reference is [66].

Some states which satisfy the Astekhar constraints are given by the loop representation, which can be introduced from the construct (depending both on the gauge field A and on a parametrized loop γ)

$$W(\gamma, A) \equiv tr P e^{\oint_\gamma A} \quad (50)$$

and a functional transform mapping functionals of the gauge field $\psi(A)$ into functionals of loops, $\psi(\gamma)$:

$$\psi(\gamma) \equiv \int \mathcal{D}A W(\gamma, A) \psi(A) \quad (51)$$

When one divides by diffeomorphisms, it is found that functions of knot classes (diffeomorphisms classes of smooth, non self-intersecting loops) satisfy all the constraints.

Some particular states sought to reproduce smooth spaces at coarse graining are the *Weaves*. It is not clear to what extent they also approach the conjugate variables (that is, the extrinsic curvature) as well.

In the presence of a cosmological constant the hamiltonian constraint reads:

$$\epsilon_{ijk} E^{ai} E^{bj} (F_{ab}^k + \frac{\lambda}{3} \epsilon_{abc} E^{ck}) = 0 \quad (52)$$

A particular class of solutions of the constraint [60] are self-dual solutions of the form

$$F_{ab}^i = -\frac{\lambda}{3} \epsilon_{abc} E^{ci} \quad (53)$$

Kodama ([41]) has shown that the Chern-Simons state

$$\psi_{CS}(A) \equiv e^{\frac{3}{2\lambda} S_{CS}(A)} \quad (54)$$

is a solution of the hamiltonian constraint. He even suggested that the *sign* of the coarse grained, classical cosmological constant was always positive, irrespectively of the sign of the quantum parameter λ , but it is not clear whether this result is general enough. There is some concern [71] that this state as such is not normalizable with the usual norm. It has been argued that this is only natural, because the physical relevant norm must be very different from the naïve one (cf. [59]) and indeed normalizability of the Kodama state has been suggested as a criterion for the correctness of the physical scalar product (cf. for example the discussion in [24]) or else that a euclidean interpretation could be given to it.

Loop states in general (suitable symmetrized) can be represented as spin network ([56]) states: colored lines (carrying some $SU(2)$ representation) meeting at nodes where intertwining $SU(2)$ operators act. A beautiful graphical representation of the group theory has been succesfully adapted for this purpose. There is a clear relationship between this representation and the Turaev-Viro [68] invariants. Many of these ideas have been foresighted by Penrose (cf. [48]).

There is also a path integral representation, known as *spin foam* (cf.[12]), a topological theory of colored surfaces representing the evolution of a spin network. These are closely related to topological BF theories, and many independent generalizations have been proposed. Spin foams can also be considered as an independent approach to the quantization of the gravitational field.([14])

In addition to its specific problems, this approach shares with all canonical approaches to covariant systems the problem of time. It is not clear its definition, at least in the absence of matter. Dynamics remains somewhat mysterious; the hamiltonian constraint does not say in what sense (with respect to what) the three-dimensional dynamics evolve.

4.1 Big results of this approach.

One of the main successes of the loop approach is that the area (as well as the volume) operator is discrete. This allows, assuming that a black hole has been formed (which is a process that no one knows how to represent in this setting), to explain the formula for the black hole entropy . The result is expressed in terms of the Barbero-Immirzi parameter ([57]). The physical meaning of this dependence is not well understood.

It has been pointed out [15] that there is a potential drawback in all theories in which the area (or mass) spectrum is discrete with eigenvalues A_n if the level spacing between eigenvalues δA_n is uniform because of the predicted thermal character of Hawking's radiation. The explicit computations in the present setting, however, lead to an space between (dimensionless) eigenvalues

$$\delta A_n \sim e^{-\sqrt{A_n}}, \quad (55)$$

which seemingly avoids this set of problems.

It has also been pointed out that [23] not only the spin foam, but almost all other theories of gravity can be expressed as topological BF theories with constraints. While this is undoubtedly an interesting and potentially useful remark, it is important to remember that the difference between the linear sigma model (a free field theory) and the nonlinear sigma models is just a matter of constraints. This is enough to produce a mass gap and asymptotic freedom in appropriate circumstances.

5 Euclidean quantum gravity

It can be boldly asserted that just by analogy with ordinary quantum field theory, the wave functional of quantum gravity must be given by:

$$\psi[h] \equiv \int_{g(\partial M)=h} \mathcal{D}g e^{-S_E[g]} \quad (56)$$

where we integrate over all riemannian metrics that obey the relevant boundary conditions, and the Einstein-Hilbert action has to be supplemented with boundary terms. This approach is problematic from the very beginning, due to the fact that the Wick analytic continuation of a lorentzian space-time is not riemannian in general (not even real), so that the whole setup seems to demand the study of real sections in a complex formulation. The point of view put forward by Hawking and collaborators [32] is that the needed analytical continuations could be hopefully made after Green's functions are evaluated.

There however is a well-known mathematical theorem of Markov (explained for physicists in [5]) asserting that there is no algorithmic way of predicting when two arbitrary manifolds are homeomorphic $M_1 \sim M_2$. The problem stems essentially from the fundamental group: any finitely presented group can be the $\pi_1(M)$ of a four-dimensional manifold, M . So one proof of the result is to simply write down a family of groups G_k such that we cannot algorithmically recognise when $G = \{e\}$. There are, in addition, further subtleties with the diffeomorphism class in $d = 4$: there is a uncountable set of non equivalent differentiable structures in $\mathbb{R}^4$: the so-called *exotic* $\mathbb{R}^4$ (cf. [18] for a physical approach; a relevant recent reference is [50]) .

Working in lorentzian signature, a Hamiltonian path integral can be dreamt of, where a functional integral is performed over three-dimensional geometries (cf. [6] only. Here the situation is slightly better: it seems that there is recent progress in the proof of Thurston's geometrization conjecture, which implies in particular Poincaré's conjecture, and which explains all three-dimensional manifolds in terms of eight different *geometries*. Incidentally, the work of Perelman ([49]) uses what mathematicians call the Ricci flow, which is *exactly*

the flow of the renormalization group of the sigma model associated to the bosonic string in a curved background.

Let us finally comment that even if the basic theory of Nature is topological one needs to *enumerate* topologies to discriminate between different ones. Besides, topological symmetry has to be broken at low energies.

In order to reach a probabilistic interpretation, a scalar product ought to be defined. The one which is naturally associated to the Wheeler-de Witt equation is not positive-definite, so this remains as an open problem in this approach.

Were somebody apply these ideas to the whole Universe (the so called Quantum Cosmology) there are other problems in store. It is not clear what is the physical interpretation of probabilities associated to a single event. A related problem is the one of the physical interpretation of Quantum Mechanics without classical observers. Many people have related this to the decoherence mechanisms (cf. for example [30]) but it seems to me that the situation is still to be clarified.

6 Perturbative (graviton) approach

A much more modest approach is to study gravitons as ordinary (massless, spin two) particles in Minkowski space-time.

$$g_{\alpha\beta} = \bar{g}_{\alpha\beta} + \kappa h_{\alpha\beta} \quad (57)$$

It seems to many people (including the author) that this is at least a preliminary step before embarking in more complicated adventures. As a quantum field theory, quantum general relativity has got a dimensionful coupling : $d(\kappa) = -1$, which means that it is not renormalizable in the usual sense of the word.

In spite of this, the theory is one loop finite on shell , as was shown in a brilliant calculation by G. 't Hooft and M. Veltman ([33]). They computed the counterterm:

$$\Delta L^{(1)} = \frac{\sqrt{\bar{g}}}{\epsilon} \frac{203}{80} \bar{R}^2 \quad (58)$$

No more miracles are expected for higher loops, and none happen. Goroff and Sagnotti ([27]) were the first to show that to two loops,

$$\Delta L^{(2)} = \frac{209}{2880(4\pi)^4} \frac{1}{\epsilon} \bar{R}^{\alpha\beta}_{\gamma\delta} \bar{R}^{\gamma\delta}_{\rho\sigma} \bar{R}^{\rho\sigma}_{\alpha\beta} \quad (59)$$

The general structure of perturbation theory is governed by the fact we have just mentioned that the coupling constant is dimensionful. A general diagram will then behave in the s-channel as $\kappa^n s^n$ and counterterms as:

$$\Delta L \sim \sum \int \kappa^n R^{(2+n/2)} \quad (60)$$

(where a symbolic notation has been used, packing all invariants with the same dimension; for example, R^2 stands for an arbitrary combination of R^2 , $R_{\alpha\beta}R^{\alpha\beta}$ and $R_{\alpha\beta\gamma\delta}R^{\alpha\beta\gamma\delta}$) conveying the fact that the theory is non-renormalizable.

It may however be pondered whether effective lagrangians are really useful for $E \ll m_P$. This possibility has been forcefully explored by Donoghue and collaborators (cf. [21]). There are some caveats: for example, when horizons are present, it seems necessary in order to be able to apply these ideas, to use some particular foliations, the so called *nice slices*). The mere fact that we are unable to predict the cosmological constant (which is the mother of all infrared problems) means that our understanding has ample room for improvement.

Could it be that in spite of the fact that general relativity is not renormalizable, there is a non perturbative sector in which the theory makes sense as a quantum theory? First of all, were that true, it would be most remarkable: there are no known QFT which are defined only nonperturbatively. Besides, at the classical level, perturbation theory works wonderfully, and there is indeed a whole framework, the parametrized post-newtonian (PPN) formalism to discriminate between alternate theories of gravity. It is then most unclear why at the quantum (and only there) level perturbation theory should fail.

We want to mention in closing this chapter, some fascinating relationships uncovered by Z. Bern and collaborators (cf. [16]) between purely field-theoretical S matrix elements

in (super)gravity and gauge theories: the so called *Gravity=Gauge* $\times$ *Gauge* conjecture. In spite of several attempts, it is not clear how this can be understood from the Einstein-Hilbert action. The relationship is of course automatic in strings, because closed string amplitudes (which include the graviton) are given by products of open string amplitudes (which contain the gauge fields). The KLT relations ([40]) are a quantitative formulation of this fact.

In view of all this, one can try to study particular extensions of the Einstein-Hilbert actions. Modifications quadratic in the curvature improve renormalizability ([39]), but have problems with unitarity at a very fundamental level ([61]).

Local supersymmetry is expected to improve the ultraviolet behavior through cancellations between fermionic and bosonic degrees of freedom. In spite of that, some infinities are allowed by the symmetries of the problem, and are thus expected to appear; for example in extended supergravities this is expected to happen at loop order $L > \frac{10}{D-2}$ in the maximally supersymmetric case in which there are 32 supercharges.

The (sad) conclusion of all this is that ordinary QFT (with a finite number of fields) does not work, even for describing small (quantum) ripples in Minkowski space.

7 Strings

It should be clear by now that we probably still do not know what is exactly the problem to which string theories are the answer. At any rate, the starting point is that all elementary particles are viewed as quantized excitations of a one dimensional object, the string, which can be either open (free ends) or closed (a loop). Excellent books are available, such as [29][52].

String theories enjoy a convoluted history. Their origin can be traced to the Veneziano model of strong interactions. A crucial step was the reinterpretation by Scherk and Schwarz ([58]) of the massless spin two state in the closed sector (previously thought to be related

to the Pomeron) as the graviton and consequently of the whole string theory as a potential theory of quantum gravity, and potential unified theories of all interactions. Now the wheel has made a complete turn, and we are perhaps back through the Maldacena conjecture ([44]) to a closer relationship than previously thought with ordinary gauge theories.



Figure 3: String theorist at work.

From a certain point of view, their dynamics is determined by a two-dimensional non-linear sigma model, which geometrically is a theory of imbeddings of a two-dimensional surface (the world sheet of the string) to a (usually ten-dimensional) target space:

$$x^\mu(\xi) : \Sigma_2 \rightarrow M_n \quad (61)$$

There are two types of interactions to consider. Sigma model interactions (in a given two-dimensional surface) are defined as an expansion in powers of momentum, where a new dimensionful parameter, $\alpha' \equiv l_s^2$ sets the scale. This scale is *a priori* believed to be of the order of the Planck length. The first terms in the action always include a coupling to the

massless backgrounds: the spacetime metric, the two-index Maxwell like field known as the Kalb-Ramond or b -field, and the dilaton. To be specific,

$$S = \frac{1}{l_s^2} \int_{\Sigma_2} g_{\mu\nu}(x(\xi)) \partial_a x^\mu(\xi) \partial_b x^\nu(\xi) \gamma^{ab}(\xi) + \dots \quad (62)$$

There are also string interactions, (changing the two-dimensional surface) proportional to the string coupling constant, g_s , whose variations are related to the logarithmic variations of the dilaton field. Open strings (which have gluons in their spectrum) *always* contain closed strings (which have gravitons in their spectrum) as intermediate states in higher string order (g_s) corrections. This interplay open/closed is one of the most fascinating aspects of the whole string theory.

It has been discovered by Friedan (cf. [25]) that in order for the quantum theory to be consistent with all classical symmetries (diffeomorphisms and conformal invariance), the beta function of the generalized couplings ⁵ must vanish:

$$\beta(g_{\mu\nu}) = R_{\mu\nu} = 0 \quad (63)$$

This result remains until now as one of the most important ones in string theory, hinting at a deep relationship between Einstein's equations and the renormalization group.

Polyakov ([53]) introduced the so called *non-critical strings* which have in general a two-dimensional cosmological constant (forbidden otherwise by Weyl invariance). The dynamics of the conformal mode (often called Liouville in this context) is, however, poorly understood.

Fundamental strings live in D=10 spacetime dimensions, and so a Kaluza-Klein mechanism of sorts must be at work in order to explain why we only see four non-compact dimensions at low energies. Strings have in general tachyons in their spectrum, and the only way to construct seemingly consistent string theories (cf. [26]) is to project out those states,

⁵There are corrections coming from both dilaton and Kalb-Ramond fields. The quoted result is the first term in an expansion in derivatives, with expansion parameter $\alpha' \equiv l_s^2$.

which leads to supersymmetry. This means in turn that all low energy predictions heavily depend on the supersymmetry breaking mechanisms.

String perturbation theory is probably well defined although a full proof is not available.

Several stringy symmetries are believed to be exact: T-duality, relating large and small compactification volumes, and S -duality, relating the strong coupling regime with the weak coupling one. Besides, extended configurations (D branes); topological defects in which open strings can end are known to be important [51]. They couple to Maxwell-like fields which are p-forms called Ramond-Ramond (RR) fields. These dualities [36] relate all five string theories (namely, Heterotic $E(8) \times E(8)$, Heterotic $SO(32)$, Type I , IIA and IIB) and it is conjectured that there is an unified eleven -dimensional theory, dubbed M -theory of which $\mathcal{N} = 1$ supergravity in $d = 11$ dimensions is the low energy limit.

7.1 Big results

Perhaps the main result is that graviton physics in flat space is well defined for the first time, and this is no minor accomplishment.

Besides, there is evidence that at least some geometric singularities are harmless in the sense that strings do not feel them. Topology change amplitudes do not vanish in string theory.

The other Big Result [62] is that one can correctly count states of extremal black holes as a function of charges. This is at the same time astonishing and disappointing. It clearly depends strongly on the objects being BPS states (that is, on supersymmetry), and the result has not been extended to non-supersymmetric configurations. On the other hand, as we have said, it *exactly* reproduces the entropy as a function of a sometimes large number of charges, without any adjustable parameter.

7.2 The Maldacena conjecture

Maldacena [44] proposed as a conjecture that *IIB* string theories in a background $AdS_5 \times S_5$ with common radius $l \sim l_s(g_s N)^{1/4}$ and N units of RR flux that is, $\int_{S_5} F_5 = N$ (which implies that $F_5 \sim \frac{N}{r^5}$) is equivalent to a four dimensional ordinary gauge theory in flat four-dimensional Minkowski space, namely $\mathcal{N} = 4$ super Yang-Mills with gauge group $SU(N)$ and coupling constant $g = g_s^{1/2}$.

Although there is much supersymmetry in the problem and the kinematics largely determine correlators, (in particular, the symmetry group $SO(2, 4) \times SO(6)$ is realized as an isometry group on the gravity side and as an R -symmetry group as well as conformal invariance on the gauge theory side) this is not fully so ⁶ and the conjecture has passed many tests in the semiclassical approximation to string theory.

The action of the RR field, given schematically by $\int F_5^2$, scales as N^2 , whereas the ten-dimensional Einstein-Hilbert $\int R$, depends on the overall geometric scale as the eighth power of the common radius, l^8 . The 't Hooft coupling is $\lambda = g^2 N \sim \frac{l^4}{l_s^4}$ and the tenth dimensional Newton's constant is $\kappa_{10}^2 \sim G_{10} \sim l_p^8 = g_s^2 l_s^8 \sim \frac{l^8}{N^2}$.

If we consider the effective five dimensional theory after compactifying on a five sphere of radius r , the RR term yields a negative contribution $\sim -(\frac{N}{r^5})^2 r^5$, whereas the positive curvature of the five sphere S^5 gives a positive contribution, $\sim \frac{1}{r^2} r^5$. The competition between these two terms in the effective potential is responsible for the minimum with negative cosmological constant.

The way the dictionary works in detail [69] is that the supergravity action corresponding to fields with prescribed boundary values is related to gauge theory correlators of certain gauge invariant operators corresponding to the particular field studied:

$$\left. e^{-S_{sugra}[\Phi_i]} \right|_{\Phi_i|_{\partial AdS} = \phi_i} = \langle e^{\int \mathcal{O}_i \phi_i} \rangle_{CFT} \quad (64)$$

⁶The only correlators that are completely determined through symmetry are the two and three-point functions.

This is the first time that a precise holographic description of spacetime in terms of a (boundary) gauge theory is proposed and, as such it is of enormous potential interest. It has been conjectured by 't Hooft [64] and further developed by Susskind [63] that there should be much fewer degrees of freedom in quantum gravity than previously thought. The conjecture claims that it should be enough with one degree of freedom per unit Planck surface in the two-dimensional boundary of the three-dimensional volume under study. The reason for that stems from an analysis of the Bekenstein-Hawking [15][31] entropy associated to a black hole, given in terms of the two-dimensional area A ⁷ of the horizon by

$$S = \frac{c^3}{4G\hbar} A. \quad (66)$$

This is a deep result indeed, still not fully understood.

It is true on the other hand that the Maldacena conjecture has only been checked for the time being in some corners of parameter space, namely when strings can be approximated by supergravity in the appropriate background.




Figure 4: Conjectured relationships between string theories in different dimensions.

⁷The area of the horizon for a Schwarzschild black hole is given by:

$$A = \frac{8\pi G^2}{c^4} M^2 \quad (65)$$

8 Dualities and branes

The so- called *T-duality* is the simplest of all dualities and the only one which can be shown to be true, at least in some contexts. At the same time it is a very stringy characteristic, and depends in an essential way on strings being extended objects. In a sense, the web of dualities rests on this foundation, so that it is important to understand clearly the basic physics involved. Let us consider strings living on an external space with one compact dimension, which we shall call y , with topology S^1 and radius R . The corresponding field in the imbedding of the string, which we shall call y (*i.e.* we are dividing the target-spacetime dimensions as (x^μ, y) , where y parametrizes the circle), has then the possibility of winding around it:

$$y(\sigma + 2\pi, \tau) = y(\sigma, \tau) + 2\pi Rm . \quad (67)$$

A closed string can close in general up to an isometry of the external spacetime.

The zero mode expansion of this coordinate (that is, forgetting about oscillators) would then be

$$y = y_c + 2p_c\tau + mR\sigma . \quad (68)$$

Canonical quantization leads to $[y_c, p_c] = i$, and single-valuedness of the plane wave $e^{iy_cp_c}$ enforces as usual $p_c \in \mathbb{Z}/R$, so that $p_c = \frac{n}{R}$.

The zero mode expansion can then be organized into left and right movers in the following way

$$\begin{aligned} y_L(\tau + \sigma) &= y_c/2 + \left(\frac{n}{R} + \frac{mR}{2} \right) (\tau + \sigma) , \\ y_R(\tau - \sigma) &= y_c/2 + \left(\frac{n}{R} - \frac{mR}{2} \right) (\tau - \sigma) . \end{aligned} \quad (69)$$

The mass shell conditions reduce to

$$\begin{aligned} m_L^2 &= \frac{1}{2} \left(\frac{n}{R} + \frac{mR}{2} \right)^2 + N_L - 1 , \\ m_R^2 &= \frac{1}{2} \left(\frac{n}{R} - \frac{mR}{2} \right)^2 + N_R - 1 . \end{aligned} \quad (70)$$

Level matching, $m_L = m_R$, implies that there is a relationship between momentum and winding numbers on the one hand, and the oscillator excess on the other

$$N_R - N_L = nm . \quad (71)$$

At this point it is already evident that the mass formula is invariant under

$$R \rightarrow R^* \equiv 2/R , \quad (72)$$

and exchanging momentum and winding numbers. This is the simplest instance of *T-Duality*.

On the other hand, it is an old observation (which apparently originated in Schrödinger) that Maxwell's equations are almost symmetrical with respect to interchange between electric and magnetic degrees of freedom. This idea was explored by Dirac and eventually lead to the discovery of the consistency conditions that have to be fulfilled if there are magnetic monopoles in nature. The fact that nonsingular magnetic monopoles appear as classical solutions in some gauge theories led further support to this duality viewpoint. In order to be able to make a consisting conjecture, first put forward by Montonen and Olive [45], supersymmetry is needed, as first remarked by Osborn [47].

Now in strings there are the so- called Ramond-Ramond (RR) fields, which are p- forms of different degrees. In the same way that one forms (i.e., the Maxwell field) couples to charged particles that is, from the spacetime point of view, to objects of dimension 0 with one-dimensional trajectories, a p-form

$$A_{\mu_1 \dots \mu_p} \quad (73)$$

would couple to a $(p - 1)$ -dimensional object, whose world history is described by a p -dimensional hypersurface

$$x^\mu = x^\mu(\xi_1 \dots \xi_p) \quad (74)$$

These objects are traditionally denoted by the name p -branes (it all originated in a dubious

joke). That is, ordinary particles are 0-branes, a string is a 1-brane, a membrane is a 2-brane, and so on.

Dualities relate branes of different dimensions in different theories; this means that if one is to take this symmetry seriously, it is not clear at all that strings are the more fundamental objects: in the so called M -theory branes appear as fundamental as strings.

If we are willing to make the hypothesis that supersymmetry is not going to be broken whilst increasing the coupling constant, g_s , some astonishing conclusions can be drawn. Assuming this, massless quanta can become massive as g_s grows only if their number, charges and spins are such that they can combine into massive multiplets (which are all larger than the irreducible massless ones). The only remaining issue, then, is whether any other massless quanta can appear at strong coupling.

Now, in the IIA string theory there are states associated to the Ramond-Ramond (RR) one form, A_1 , namely the D-0-branes, whose tension goes as $m \sim \frac{1}{g_s}$. This clearly gives new massless states in the strong coupling limit.

There are reasons⁸ to think that this new massless states are the first level of a Kaluza-Klein tower associated to compactification on a circle of an 11-dimensional theory. Actually, assuming an 11-dimensional spacetime with an isometry $k = \frac{\partial}{\partial y}$, an Ansatz which exactly reproduces the dilaton factors of the IIA string is

$$ds_{(11)}^2 = e^{\frac{4}{3}\phi} (dy - A_\mu^{(1)} dx^\mu)^2 + e^{-\frac{2}{3}\phi} g_{\mu\nu} dx^\mu dx^\nu . \quad (75)$$

Equating the two expressions for the D0 mass,

$$\frac{1}{g_s} = \frac{1}{R_{11}} , \quad (76)$$

leads to $R_{11} = e^{\frac{2}{3}\phi} = g_A^{2/3}$.

This means that a new dimension appears at strong coupling, and this dimension is related to the dilaton. The only reason why we do not see it at low energiew is precisely

⁸In particular: The fact that there is the possibility of a central extension in the IIA algebra, related to the Kaluza-Klein compactification of the d=11 Supergravity algebra.

because of the smallness of the string coupling, related directly to the dilaton field. The other side of this is that this eleven dimensional theory, dubbed *M-theory* does not have any weak coupling limit; it is always strongly coupled. Consequently, not much is known on this theory, except for the fact that its field theory, low curvature limit is $\mathcal{N} = 1$ supergravity in $d = 11$ dimensions.

All supermultiplets of massive one-particle states of the IIB string supersymmetry algebra contain states of at least spin 4. This means that under the previous set of hypothesis, the set of massless states at weak coupling must be exactly the same as the corresponding set at strong coupling. This means that there must be a symmetry mapping weak coupling into strong coupling.

There is a well-known candidate for this symmetry: Let us call, as usual, l the RR scalar and ϕ the dilaton (NSNS). We can pack them together into complex scalar

$$S = l + ie^{-\frac{\phi}{2}} . \quad (77)$$

The IIB supergravity action in d=10 is invariant under the $SL(2, \mathbb{R})$ transformations

$$S \rightarrow \frac{aS + b}{cS + d} , \quad (78)$$

if at the same time the two two-forms, $B_{\mu\nu}$ (the usual, ever-present, NS field), and $A^{(2)}$, the RR field transform as

$$\begin{pmatrix} B \\ A^{(2)} \end{pmatrix} \rightarrow \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} B \\ A^{(2)} \end{pmatrix} , \quad (79)$$

Both the, Einstein frame, metric $g_{\mu\nu}$ and the four-form $A^{(4)}$ are inert under this $SL(2, \mathbb{R})$ transformation.

A discrete subgroup $SL(2, \mathbb{Z})$ of the full classical $SL(2, \mathbb{R})$ is believed to be an exact symmetry of the full string theory. The exact imbedding of the discrete subgroup in the full $SL(2, \mathbb{R})$ depends on the vacuum expectation value of the RR scalar.

The particular transformation

$$g = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} , \quad (80)$$

maps ϕ into $-\phi$ (when $l = 0$), and B into $A^{(2)}$. This means that the string coupling

$$g_s \rightarrow \frac{1}{g_s} \quad (81)$$

This is a strong/weak coupling type of duality, similar to the electromagnetic duality in that sense. The standard name for it is an *S-duality* type of transformation, mapping the ordinary string with NS charge, to another string with RR charge (which then must be a D-1-brane, and is correspondingly called a *D-string*), and, from there, is connected to all other D-branes by T-duality.

Using the fact that upon compactification on S^1 , IIA at R_A is equivalent to IIB at $R_B \equiv 1/R_A$, and the fact that the effective action carries a factor of $e^{-2\phi}$ we get

$$R_A g_B^2 = R_B g_A^2 , \quad (82)$$

which combined with our previous result, $g_A = R_{11}^{3/2}$ implies that $g_B = \frac{R_{11}^{3/2}}{R_A}$. Now the Kaluza-Klein Ansatz implies that from the eleven dimensional viewpoint the compactification radius is measured as

$$R_{10}^2 \equiv R_A^2 e^{-2\phi/3} , \quad (83)$$

yielding

$$g_B = \frac{R_{11}}{R_{10}} . \quad (84)$$

From the effective actions written above it is easy to check that there is a (S-duality type) field transformation mapping the SO(32) Type I open string into the SO(32) Heterotic one namely

$$\begin{aligned} g_{\mu\nu} &\rightarrow e^{-\phi} g_{\mu\nu}^{Het} , \\ \phi &\rightarrow -\phi , \\ B' &\rightarrow B . \end{aligned} \quad (85)$$

This means that physically there is a strong/weak coupling duality, because coupling constants of the compactified theories would be related by

$$\begin{aligned} g_{het} &= 1/g_I , \\ R_{het} &= R_I/g_I^{1/2} . \end{aligned} \tag{86}$$

9 Summary: the state of the art in quantum gravity

In the loop approach one is working with nice candidates for a quantum theory. The theories are interesting, probably related to topological field theories ([17]) and background independence as well as diffeomorphism invariance are clearly implemented. On the other hand, it is not clear that their low energy limit is related to Einstein gravity.

Strings start from a perturbative approach more familiar to a particle physicist. However, they carry all the burden of supersymmetry and Kaluza-Klein. It has proved to be very difficult to study nontrivial non-supersymmetric dynamics.

Finally, and this applies to all approaches, the holographic ideas seem intriguing; there are many indications of a deep relationship between gravity and gauge theories.

We would like to conclude by insisting on the fact that although there is not much we know for sure on quantum effects on the gravitational field, even the few things we know are a big feat, given the difficulty to do physics without experiments.

Progress could be made if we could derive semiclassical gravity in such a way that corrections to it can be reliably estimated, for example

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 2\kappa^2 \langle \psi | \hat{T}_{\mu\nu} | \psi \rangle + \frac{1}{L^2}\Delta, \tag{87}$$

when working at a certain scale of distances, say L . In order to understand those equations, we would had to know something about the operator of which the first member is the expectation value; something about the state on which the expectation value is computed (In particular, if it is the vacuum, how is it to be defined?) and finally, something about

the definition of the energy momentum tensor as a composite operator. A question of obvious physical interest is the estimate of the size of the corrections: Is the expected error at a given scale of distance L

$$\Delta \sim \frac{\hbar G}{c^3 L^2} \quad (88)$$

or, does it depend of the characteristic energy of the source?

$$\Delta \sim \frac{GE^2}{\hbar c^5} \quad (89)$$

measured with respect to what?

It is painfully clear that there is still a large margin for improving our understanding of effective quantum field theories. For example, there is still no convincing derivation of Hawking radiation without transplanckian modes appearing at some point (this particular example is related to the existence of the nice slices mentioned above). Besides, we do not understand the cosmological constant, which is clearly related to the estimate of Δ .

The observational prospects are rather poor. In many models, in particular in the loop approach (and also in strings, with some qualifications) deviations from the lorentzian dispersion relations are expected:

$$E^2 = \vec{p}^2 + m^2 + E^2 \sum_{n=1} c_n \left(\frac{E}{m_P} \right)^n \quad (90)$$

Other contributions will undoubtedly analyze those in much more detail. Let us now simply mention that noncommutative models make similar predictions.

Winding states are stringy phenomena, and its observation would be very interesting. Stringy predictions, however, are in general difficult to disentangle from predictions of supersymmetry (SUSY). Namely, SUSY has to be broken, and this scale spoils almost all differences between strings and QFT models.

With the great triumph of particle physics at the end of the seventies, namely the *experimental* discovery of the intermediate bosons related to electroweak interactions, the standard model was confirmed in all its essential traits, waiting only for the Higgs to

be discovered (at LHC?) and the theoretical effort has concentrated in more and more speculative topics, and experimental guidance has become correspondingly scarce. The net result is that, even more so that in the old days of the hunting for the theory of strong interactions, theoretical physics is divided in almost disconnected clans.

All this is even more true when talking about quantum gravity, a paradise of speculation.

This is the reason why all efforts such as the one in the present workshop, aiming at making contact with experiment and/or observation are welcome, and will eventually redirect physics on a healthier track when we learn to recognise the physically relevant facts that presumably lie in front of our eyes.

10 Acknowledgements

I am grateful to Giovanni Amelino-Camelia and to Jerzy Kowalski-Glikman for organizing such a stimulating workshop and for allowing me to take part in it. I am also grateful to Carlos Muñoz for lending his picture of a stringer . Finally, I am indebted to Jorge Conde for reading the manuscript.

References

- [1] R. Arnowitt, S. Deser, and C. W. Misner, “ Canonical Variables for General Relativity” Phys. Rev. 117, 1595-1602 (1960)
- [2] J. Alfaro and G. Palma, “Loop quantum gravity corrections and cosmic rays decays,” Phys. Rev. D 65 (2002) 103516 [arXiv:hep-th/0111176].
“Loop quantum gravity and ultra high energy cosmic rays,” Phys. Rev. D 67 (2003) 083003 [arXiv:hep-th/0208193].

- [3] E. Alvarez, “Quantum Gravity: A Pedagogical Introduction To Some Recent Results,”
Rev. Mod. Phys. 61 (1989) 561.
- [4] E. Alvarez, “Low-Energy Effects Of Quantum Gravity,” FTUAM-89-15 *Lectures given
at Mtg. on Recent Developments in Gravitation, Barcelona, Spain, Sep 5-8, 1989*
- [5] E. Alvarez, “Some general problems in quantum gravity,” CERN-TH-6257-91 *Lec-
tures at the 22nd Gift Int. Seminar in Theoretical Physics: Quantum Gravity and
Cosmology, S. Feliu De Guixols, Jun 3-6, 1991*
- [6] E. Alvarez, “Some general problems in quantum gravity. 2. The Three-dimensional
case,” Int. J. Mod. Phys. D 2 (1993) 1 [arXiv:hep-th/9211050].
- [7] E. Alvarez, L. Alvarez-Gaume and Y. Lozano, “An introduction to T duality in string
theory,” Nucl. Phys. Proc. Suppl. 41 (1995) 1 [arXiv:hep-th/9410237].
- [8] L. Alvarez-Gaume and M. A. Vazquez-Mozo, “Topics in string theory and quantum
gravity,” arXiv:hep-th/9212006.
- [9] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, “New dimensions
at a millimeter to a Fermi and superstrings at a TeV,” Phys. Lett. B 436 (1998) 257
[arXiv:hep-ph/9804398].
- [10] A. Ashtekar, “New Hamiltonian Formulation Of General Relativity,” Phys. Rev. D
36 (1987) 1587.
- [11] A. Ashtekar, “Quantum Geometry And Black Hole Entropy”, gr-qc/0005126
[=Adv.Theor.Math.Phys. 4 (2000) 1]
“Quantum Geometry Of Isolated Horizons And Black Hole Entropy”, Phys.Rev. D57
(1998) 1009 [=gr-qc/9705059]
- [12] J.Baez, “An Introduction To Spin Foam Models Of Quantum Gravity And Bf Theory”,
gr-qc/0010050

- [13] J. F. Barbero, “Real Ashtekar variables for Lorentzian signature space times,” Phys. Rev. D 51 (1995) 5507 [arXiv:gr-qc/9410014].
- [14] J. W. Barrett and L. Crane, “A Lorentzian signature model for quantum general relativity,” Class. Quant. Grav. 17 (2000) 3101 [arXiv:gr-qc/9904025].
- [15] J. Bekenstein, *Black Holes And Entropy* Phys.Rev. D7 (1973) 2333
(with V. Mukhanov), “Spectroscopy Of The Quantum Black Hole”, Commun.Math.Phys. 125 (1989) 417
- [16] Z. Bern, “Perturbative quantum gravity and its relation to gauge theory,” Living Rev. Rel. 5 (2002) 5 [arXiv:gr-qc/0206071].
- [17] M. Blau, “Topological Gauge Theories Of Antisymmetric Tensor Fields”, hep-th/9901069 [=Adv.Theor.Math.Phys. 3 (1999) 1289]
- [18] C. H. Brans, “Exotic smoothness structures in physics,” *Prepared for 1st Mexican School on Gravitation and Mathematical Physics, Guanajuato, Mexico, 12-16 Dec 1994*
- [19] S. R. Coleman and S. L. Glashow, “High-energy tests of Lorentz invariance,” Phys. Rev. D 59 (1999) 116008 [arXiv:hep-ph/9812418].
- [20] B. S. Dewitt, “Quantum Theory Of Gravity. 1. The Canonical Theory,” Phys. Rev. **160** (1967) 1113.
“Quantum Theory of Gravity. II. The Manifestly Covariant Theory”, Phys. Rev. 162, 1195-1239 (1967).
- [21] J. F. Donoghue, “General Relativity As An Effective Field Theory: The Leading Quantum Corrections,” Phys. Rev. D **50** (1994) 3874 [arXiv:gr-qc/9405057].

- [22] M. J. Duff, “Inconsistency Of Quantum Field Theory In Curved Space-Time,”
ICTP/79-80/38 *Talk presented at 2nd Oxford Conf. on Quantum Gravity, Oxford, Eng., Apr 1980*
- [23] L. Freidel, K. Krasnov and R. Puzio, “BF description of higher-dimensional gravity theories,” *Adv. Theor. Math. Phys.* 3 (1999) 1289 [arXiv:hep-th/9901069].
- [24] L. Freidel and L. Smolin, “The linearization of the Kodama state,”
arXiv:hep-th/0310224.
- [25] D. Friedan, “Nonlinear Models In Two Epsilon Dimensions,” *Phys. Rev. Lett.* 45 (1980) 1057.
- [26] F. Gliozzi, J. Scherk and D. I. Olive, “Supersymmetry, Supergravity Theories And The Dual Spinor Model,” *Nucl. Phys. B* 122 (1977) 253.
- [27] M. H. Goroff and A. Sagnotti, “The Ultraviolet Behavior Of Einstein Gravity,” *Nucl. Phys. B* **266**, 709 (1986).
- [28] M. B. Green and J. H. Schwarz, “Anomaly Cancellation In Supersymmetric D=10 Gauge Theory And Superstring Theory,” *Phys. Lett. B* 149 (1984) 117.
- [29] M. B. Green, J. H. Schwarz and E. Witten, “Superstring Theory. Vol. 1: Introduction,”
“Superstring Theory. Vol. 2: Loop Amplitudes, Anomalies And Phenomenology,”
- [30] J. B. Hartle, “Space-time quantum mechanics and the quantum mechanics of space-time,” arXiv:gr-qc/9304006.
- [31] S. Hawking, *Particle Creation By Black Holes* *Commun.Math.Phys.* 43 (1975) 199
- [32] G. W. . Gibbons and S. W. . Hawking, “Euclidean Quantum Gravity,”
- [33] G. ’t Hooft and M. J. G. Veltman, “One Loop Divergencies In The Theory Of Gravitation,” *Annales Poincare Phys. Theor. A* **20** (1974) 69.

- [34] G. T. Horowitz, “Quantum gravity at the turn of the millennium,” arXiv:gr-qc/0011089.
- [35] G. Horowitz, “Exactly Soluble Diffeomorphism Invariant Theories”, *Annals Phys.* 205 (1991) 130
- [36] C. M. Hull and P. K. Townsend, “Unity of superstring dualities,” *Nucl. Phys. B* 438 (1995) 109 [arXiv:hep-th/9410167].
- [37] G. Immirzi, “Quantum gravity and Regge calculus,” *Nucl. Phys. Proc. Suppl.* 57 (1997) 65 [arXiv:gr-qc/9701052].
- [38] C. J. Isham, “Structural issues in quantum gravity,” arXiv:gr-qc/9510063.
- [39] J. Julve and M. Tonin, “Quantum Gravity With Higher Derivative Terms,” *Nuovo Cim. B* **46** (1978) 137.
- [40] H. Kawai, D. C. Lewellen and S. H. H. Tye, “A Relation Between Tree Amplitudes Of Closed And Open Strings,” *Nucl. Phys. B* **269** (1986) 1.
- [41] H. Kodama, “Holomorphic Wave Function Of The Universe,” *Phys. Rev. D* 42 (1990) 2548.
- [42] M. Luscher, R. Narayanan, P. Weisz and U. Wolff, “The Schrodinger functional: A Renormalizable probe for nonAbelian gauge theories,” *Nucl. Phys. B* 384 (1992) 168 [arXiv:hep-lat/9207009].
- [43] Y. M. Makeenko and A. A. Migdal, “Exact Equation For The Loop Average In Multicolor QCD,” *Phys. Lett. B* 88 (1979) 135 [Erratum-ibid. B 89 (1980) 437].
- [44] J. Maldacena, *The large N limit of superconformal field theories and supergravity*, *Adv.Theor.Math.Phys.*2:231-252,1998, hep-th/9711200.

- [45] C. Montonen and D. I. Olive, “Magnetic Monopoles As Gauge Particles?,” Phys. Lett. B **72** (1977) 117.
- [46] R. C. Myers and M. Pospelov, “Experimental challenges for quantum gravity,” arXiv:hep-ph/0301124.
- [47] H. Osborn, “Topological Charges For N=4 Supersymmetric Gauge Theories And Monopoles Of Spin 1,” Phys. Lett. B **83** (1979) 321.
- [48] R. Penrose, *Angular momentum: an approach to combinatorial spacetime*, in *Quantum theory and beyond*, T. Bastin ed. (Cambridge University Press, 1971)
- [49] G. Perelman, “The entropy formula for the Ricci flow and its geometric applications”
- [50] H. Pfeiffer, arXiv:gr-qc/0404088.
- [51] J. Polchinski, “Dirichlet-Branes and Ramond-Ramond Charges,” Phys. Rev. Lett. **75** (1995) 4724 [arXiv:hep-th/9510017].
- [52] J. Polchinski, “String Theory. Vol. 1: An Introduction To The Bosonic String,”
“String Theory. Vol. 2: Superstring Theory And Beyond,”
- [53] A. M. Polyakov, “Quantum Geometry Of Bosonic Strings,” Phys. Lett. B **103** (1981) 207.
- [54] C. Rovelli, “Loop quantum gravity,” Living Rev. Rel. **1** (1998) 1 [arXiv:gr-qc/9710008].
- [55] C. Rovelli *Loop Quantum Gravity* gr-qc/9606088 [=Phys.Lett. B380 (1996) 257]
“The Immirzi Parameter In Quantum General Relativity”, gr-qc/9505012
[=Phys.Lett. B360 (1995) 7]
- [56] C. Rovelli and L. Smolin, “Spin networks and quantum gravity,” Phys. Rev. D **52** (1995) 5743 [arXiv:gr-qc/9505006].

- [57] C. Rovelli and L. Smolin, “Discreteness of area and volume in quantum gravity,” Nucl. Phys. B 442 (1995) 593 [Erratum-ibid. B 456 (1995) 753] [arXiv:gr-qc/9411005].
- [58] J. Scherk and J. H. Schwarz, “Dual Models For Nonhadrons,” Nucl. Phys. B 81 (1974) 118.
- [59] L. Smolin, “How far are we from the quantum theory of gravity?,” arXiv:hep-th/0303185.
- [60] L. Smolin, “Quantum gravity with a positive cosmological constant,” arXiv:hep-th/0209079.
- [61] K. S. Stelle, “Renormalization Of Higher Derivative Quantum Gravity,” Phys. Rev. D 16 (1977) 953.
- [62] A. Strominger and C. Vafa, “Microscopic Origin of the Bekenstein-Hawking Entropy,” Phys. Lett. B 379 (1996) 99 [arXiv:hep-th/9601029].
- [63] L. Susskind, “The World as a hologram,” J. Math. Phys. 36 (1995) 6377 [arXiv:hep-th/9409089].
- [64] G. ’t Hooft, “Dimensional Reduction In Quantum Gravity,” arXiv:gr-qc/9310026.
- [65] G. ’t Hooft and M. J. Veltman, “One Loop Divergencies In The Theory Of Gravitation,” Annales Poincare Phys. Theor. A 20 (1974) 69.
- [66] T. Thiemann, “Introduction to modern canonical quantum general relativity,” arXiv:gr-qc/0110034.
- [67] T. Thiemann, “Anomaly-free formulation of non-perturbative, four-dimensional Lorentzian quantum gravity,” Phys. Lett. B 380 (1996) 257 [arXiv:gr-qc/9606088].
“Gauge field theory coherent states (GCS). I: General properties,” Class. Quant. Grav. 18 (2001) 2025 [arXiv:hep-th/0005233].

- [68] V. G. Turaev and O. Y. Viro, “State Sum Invariants Of 3 Manifolds And Quantum 6j Symbols,” *Topology* 31 (1992) 865.
- [69] E. Witten, *Anti de Sitter Space and Holography*, [hep-th/9802150](#);
Anti-de Sitter space, thermal phase transition and confinement in gauge theories
[hep-th/9803131](#).
- [70] E. Witten, “(2+1)-Dimensional Gravity As An Exactly Soluble System,” *Nucl. Phys. B* 311 (1988) 46.
- [71] E. Witten, “A note on the Chern-Simons and Kodama wavefunctions,”
[arXiv:gr-qc/0306083](#).